

Automorphisms of Projective Geometry in the sense of Klein

Navarro, A., Navarro, J.

The projective space $\mathbb{P}_n$ over a field k is usually defined as the set of 1-dimensional subspaces of a $n+1$ -dimensional k -vector space E and its automorphisms are defined to be the projectivizations of semilinear automorphisms of E (also called Staudt projectivities).

For $n \geq 2$, recall that the projective space can also be defined axiomatically as a lattice satisfying certain properties and it can be proved that automorphisms of this lattice structure (i.e., collineations) are precisely Staudt projectivities (see [1], Thm. 2.26).

On the other hand, Klein defined projective geometry as the action of the group PGL_n on the set $\mathbb{P}_n$. In this note, we prove that the automorphisms of the Klein geometry $(\mathbb{P}_n, \mathrm{PGL}_n)$ are Staudt projectivities, for any n .

Definition: A *geometry* in the sense of Klein is a pair (X, G) where X is a set and G is a subgroup of the group $\mathrm{Bij}(X)$ of all bijections $X \rightarrow X$.

Given two geometries (X, G) and (X', G') , we say that a bijection $\varphi: X \rightarrow X'$ is an *isomorphism of geometries* if the map $\Phi: \mathrm{Bij}(X) \rightarrow \mathrm{Bij}(X')$, $\Phi(g) := \varphi \circ g \circ \varphi^{-1}$, preserves the structural groups: $\Phi(G) = G'$; that is to say, $\varphi G \varphi^{-1} = G'$. Isomorphisms of geometries $\varphi: (X, G) \rightarrow (X', G')$ are said to be *automorphisms* of (X, G) .

Recall that, given a subgroup H of a group G , the *normalizer* $N(H)$ of H in G is defined to be the subgroup $N(H) = \{n \in G: nHn^{-1} = H\}$, so the group of automorphisms of (X, G) is just the normalizer of G in the group $\mathrm{Bij}(X)$.

Fundamental examples:

- The *projective geometry* $(\mathbb{P}_n, \mathrm{PGL}_n)$, where $\mathbb{P}_n$ is the set of 1-dimensional vector subspaces of a k -vector space E of dimension $n+1$ and PGL_n is the group of Poncelet projectivities (projectivizations of k -linear automorphisms of E) with the obvious action on $\mathbb{P}_n$.

- The *affine geometry* $(\mathbb{A}_n, \mathrm{Aff}_n)$, where $\mathbb{A}_n$ is the complement of an hyperplane in $\mathbb{P}_n$, named *hyperplane at infinity*, and Aff_n is the subgroup of PGL_n of all Poncelet projectivities of $\mathbb{P}_n$ preserving the hyperplane at infinity, with the action by restriction on $\mathbb{A}_n$.

- The *eclidean geometry*, $(\mathbb{A}_n(\mathbb{R}), \mathrm{Mov}_n)$, where Mov_n is the group of motions of the real affine space $\mathbb{A}_n(\mathbb{R})$.

Automorphisms of the projective geometry

Recall that a bijection $\varphi: \mathbb{P}_n \rightarrow \mathbb{P}_n$ is called a *Staudt projectivity* if it is the projectivization of a bijection $f: E \rightarrow E$ such that $f(\lambda e + \mu v) = h(\lambda)f(e) + h(\mu)f(v)$, $\forall e, v \in E$, $\lambda, \mu \in k$, for some automorphism of fields $h: k \rightarrow k$, named automorphism associated to φ .

In dimension $n \geq 2$, the so called Fundamental Theorem of Projective Geometry ([1], Thm. 2.26) states that Staudt projectivities are just collineations (i.e., bijections $\varphi: \mathbb{P}_n \rightarrow \mathbb{P}_n$ that preserve alignment). So, in order to prove Theorem 0.2, it would be useful to characterize collineations in terms of Poncelet projectivities. To do so, for any given three different points P_1, P_2, P_3 in $\mathbb{P}_n$ we consider the set:

$$H_{P_1, P_2}(P_3) = \{P \in \mathbb{P}_n: \exists \varphi \in \text{PGL}_n, \varphi(P_1) = P_1, \varphi(P_2) = P_2, \varphi(P_3) = P \neq P_3\}$$

Lemma 0.1 *Three different points $P_1, P_2, P_3 \in \mathbb{P}_n$ are collinear if and only if*

$$H_{P_1, P_2}(P_3) = H_{P_1, P_3}(P_2) = H_{P_2, P_3}(P_1)$$

Proof: If P_1, P_2, P_3 are collinear, then $H_{P_1, P_2}(P_3)$ is just the complement of P_1, P_2, P_3 in the line passing through them.

Conversely, if P_1, P_2, P_3 are not collinear, then $H_{P_1, P_2}(P_3)$ is the complement of the line R joining P_1 and P_2 (apart from P_3). Since a projective line always has more than two points, we may fix a point $Q \in R$, $Q \neq P_1, P_2$, and then it is clear that $Q \notin H_{P_1, P_2}(P_3)$ and $Q \in H_{P_1, P_3}(P_2)$. $\square$

Theorem 0.2 *The group of automorphisms of the projective geometry $(\mathbb{P}_n, \text{PGL}_n)$ is the group PSeml_n of Staudt projectivities, for any n .*

That is to say, PSeml_n is the normalizer of the group PGL_n of Poncelet projectivities in the group of all bijections of $\mathbb{P}_n$.

Proof: Let φ be a Staudt projectivity with associated automorphism $h: k \rightarrow k$. If τ is a Poncelet projectivity, then $\varphi\tau\varphi^{-1}$ is a Poncelet projectivity, as it is a Staudt projectivity with associated automorphism $h \circ \text{Id} \circ h^{-1} = \text{Id}$. Applying the same reasoning to φ^{-1} , it follows the equality, $\varphi^{-1}\text{PGL}_n\varphi = \text{PGL}_n$.

Now, let us prove that any automorphism φ of $(\mathbb{P}_n, \text{PGL}_n)$ is a Staudt projectivity.

$n > 1$. In this case Staudt projectivities are just collineations ([1], Thm 2.26) and hence it is enough to check that φ is a collineation. Since φ preserves Poncelet projectivities by hypothesis, it follows that $\varphi(H_{P_1, P_2}(P_3)) = H_{\varphi(P_1), \varphi(P_2)}(\varphi(P_3))$, and Lemma 0.1 allows to conclude that $\varphi(P_1), \varphi(P_2), \varphi(P_3)$ are collinear whenever so are P_1, P_2, P_3 .

$n = 1$. Let us fix a projective reference (P_0, P_∞, P_1) in $\mathbb{P}_1$ and write $P'_0 := \varphi(P_0)$, $P'_\infty := \varphi(P_\infty)$, $P'_1 := \varphi(P_1)$.

Consider, in the affine line $\mathbb{A}_1 = \mathbb{P}_1 - \{P_\infty\}$, the origin P_0 and the unit point P_1 , so that there is a bijection $\mathbb{A}_1 \simeq k$, and P_λ will denote the point corresponding to $\lambda \in k$ (and analogously in the affine line $\mathbb{A}'_1 = \mathbb{P}_1 - \{P'_\infty\}$).

Now the composition $k \simeq \mathbb{A}_1 \xrightarrow{\varphi} \mathbb{A}'_1 \simeq k$ defines a bijection $h: k \rightarrow k$ that is an automorphism of the field k (Lemma 0.3) and such that $\varphi(P_\lambda) = P'_{h(\lambda)}$.

Therefore, if (e_0, e_1) is a basis of E associated to the reference (P_0, P_∞, P_1) and (e'_0, e'_1) is a basis associated to the reference (P'_0, P'_∞, P'_1) , then φ is clearly the projectivization of the semilinear map $f(\lambda e_0 + \mu e_1) = h(\lambda) e'_0 + h(\mu) e'_1$. $\square$

Lemma 0.3 *The map $h: k \rightarrow k$ defined in the proof above is an automorphism of fields.*

Proof: By definition, $h(0) = 0$ and $h(1) = 1$, so we only have to show that h preserves the addition and the product. Let τ_μ and σ_μ denote the translation by $\mu \in k$ in $\mathbb{A}_1$ and the homothety with center at P_0 and ratio μ , respectively (and analogously τ'_μ and σ'_μ in $\mathbb{A}'_1$).

Firstly, let us check that $\varphi\tau_\mu\varphi^{-1}$ is $\tau'_{h(\mu)}$. since P_∞ is the unique fixed point of τ_μ , P'_∞ is the unique fixed point of $\varphi\tau_\mu\varphi^{-1}$ and hence it is a translation in $\mathbb{A}'_1$ (as it is an homography by hypothesis). Then, $\varphi\tau_\mu\varphi^{-1}$ is determined by the image of one point:

$$\varphi\tau_\mu\varphi^{-1}(P'_0) = \varphi\tau_\mu(P_0) = \varphi(P_\mu) = P'_{h(\mu)}.$$

Hence, $\varphi\tau_\mu = \tau'_{h(\mu)}\varphi$, and applying both terms to P_λ :

$$\begin{aligned}\varphi\tau_\mu(P_\lambda) &= \varphi(P_{\lambda+\mu}) = P'_{h(\lambda+\mu)} \\ \tau'_{h(\mu)}\varphi(P_\lambda) &= \tau'_{h(\mu)}(P'_{h(\lambda)}) = P'_{h(\lambda)+h(\mu)}\end{aligned}$$

it follows that $h(\lambda + \mu) = h(\lambda) + h(\mu)$.

In a similar way, it can be checked that $\varphi\sigma_\mu\varphi^{-1}$ is just the homothety $\sigma'_{h(\mu)}$, that allows to conclude that $h(\lambda\mu) = h(\lambda)h(\mu)$. $\square$

Automorphisms of the affine and euclidean geometries

In the affine case, collinear points may be characterized in terms of affinities, analogously to what is done in Lemma 0.1. Then, taking care of details, it follows:

Theorem 0.4 *The group of automorphisms of the affine geometry $(\mathbb{A}_n, \text{Aff}_n)$ over a field $k \neq \mathbb{F}_2$ is the group of (restrictions to $\mathbb{A}_n$ of) Staudt projectivities preserving the hyperplane at infinity, for any n .*

As for the euclidean case, collinear points may be characterized in terms of motions. Then, as affinities preserving motions also preserve perpendicular lines, it can be checked that:

Theorem 0.5 *For $n \geq 2$, the group of automorphisms of the Euclidean geometry $(\mathbb{A}_n(\mathbb{R}), \text{Mov}_n)$ is the group of similarities.*

Notice that the above theorem is false when $n = 1$. A counter example is any bijection $\varphi: \mathbb{R} \rightarrow \mathbb{R}$ that respects the addition, for example, any $\mathbb{Q}$ -linear automorphism of $\mathbb{R}$. If $\phi \in \text{Mov}_1$, then $\phi(x) = \pm x + b$ and it follows that $(\varphi^{-1}\phi\varphi)(x) = \varphi^{-1}(\pm\varphi(x) + b) = \pm x + \varphi^{-1}(b)$, so that φ belongs to the normalizer of Mov_1 .

However, it is not difficult to show that any *continuous* automorphism φ of the euclidean line $(\mathbb{A}_1(\mathbb{R}), \text{Mov}_1)$ is in fact a similarity $\varphi(x) = ax + b$.

An application: the outer automorphism of S_6

An application of Theorem 0.2, told to the authors by J. B. Sancho, is a construction of the outer automorphism of S_6 (for other descriptions of this exotic object see [2, 3]).

Let $k = \mathbb{Z}/5\mathbb{Z}$ be the field with 5 elements. The projective line has in this case 6 elements, so that there are $6 \cdot 5 \cdot 4$ homographies and PGL_1 is a subgroup of S_6 of index 6. Since the identity is the unique automorphism of the field k , Theorem 0.2 states that PGL_1 coincides with its normalizer in S_6 ; hence it has 6 conjugated subgroups $H_1 := \mathrm{PGL}_1, H_2, \dots, H_6$.

Now any permutation $\tau \in \mathrm{Bij}(\mathbb{P}_1) = S_6$ of the projective line defines, by conjugation, a permutation $F(\tau)$ of this set $\{H_1, \dots, H_6\}$. So we obtain an automorphism $F: S_6 \rightarrow S_6$ such that $F(\mathrm{PGL}_1)$ is contained in the subgroup of all permutations fixing the element H_1 (in fact they coincide, since both subgroups have index 6). Now, since no point of the projective line is fixed by the group of all homographies, so does its image under any inner automorphism, and we conclude that this automorphism F is an outer automorphism of S_6 .

References

- [1] Artin, E. *Geometric Algebra*, Wiley Classics Library Edition, New York (1988).
- [2] Howard, B., Millson, J., Snowden, A., Vakil, R. *A description of the outer automorphism of S_6 and the invariants of six points in projective space*, J. Combin. Theory Ser. A 115, (2008) 1296-1303
- [3] Janusz, G., Rotman, J. *Outer automorphism of S_6* , Amer. Math. Monthly, **89**, 6, (1982) 407-410